

# Cerebellar Long Term Depression as a Supervised Learning Rule with All or Nothing Character

Mitsuo Kawato

ATR Computational Neuroscience Laboratories

The cerebellar internal model theory postulates that the cerebellar cortex acquires many internal models of controlled objects, dynamical processes in the external world, and even other one's brain dependent on long-term depression (LTD) of Purkinje cells. A specific version of this theory, the feedback-error-learning model postulates that the climbing fiber inputs to Purkinje cells carry the feedback motor command, which could be regarded as an approximation to the error signal for motor commands and can supervise learning of inverse dynamics models. Many experimental supports were obtained from the ventral paraflocculus of the cerebellum during monkey control of ocular following responses. For arm movements under multiple force fields, firings of many Purkinje cells correlate with dynamics [1]. fMRI studies mapped forward and inverse models of manipulated objects and tools in the cerebellar cortex.

Kinetic models of LTD [2,3] suggest a cascade of excitable and bistable dynamical processes, which may resolve plasticity-stability dilemma at single spine level. That is, even a single pulse of climbing fiber input combined with an early train of several parallel fiber pulses can induce  $\text{Ca}^{2+}$  induced  $\text{Ca}^{2+}$  release via IP3 receptors on ER. The MAPK positive feedback loop leaky integrates resulting large  $\text{Ca}^{2+}$  elevation and if it crosses the threshold then the state moves to the depressed equilibrium. These models explain diverse LTD experiments and clearly demonstrate that LTD is a supervised learning rule, and not anti-Hebbian as erroneously characterized. The MAPK positive feedback loop model [2] was recently supported by a  $\text{Ca}^{2+}$  photo-uncaging and imaging experiment [4] that supports LTD all-or-none character.

In the LTD study, the most important information within the system  $\text{Ca}^{2+}$  can be measured and manipulated directly, thus the theory was quite rigorously proved. At the system level, the cerebellar internal models are still debated and we need to develop a method to directly manipulate information.

## References

- [1] Yamamoto K, Kawato M, Kotoaska S, Kitazawa S: Encoding of movement dynamics by Purkinje cell simple spike activity during fast arm movements under resistive and assistive force fields. *Journal of Neurophysiology*, **97**: 1588-1599, 2007.
- [2] Kuroda S, Schweighofer N, Kawato M: Exploration of signal transduction pathways in cerebellar long-term depression by kinetic simulation. *Journal of Neuroscience*, **21**: 5693-5702, 2001.
- [3] Doi T, Kuroda S, Michikawa T, Kawato M: Inositol, 1, 4, 5-trisphosphate-dependent  $\text{Ca}^{2+}$  threshold dynamics detect spike timing in cerebellar Purkinje Cells. *Journal of Neuroscience*, **25**: 950-961, 2005.
- [4] Tanaka K, Khiroug L, Santamaria F, Doi T, Ogasawara H, Ellis-Davies G, Kawato M, Augustine GJ:  $\text{Ca}^{2+}$  requirements for cerebellar long-term synaptic depression: role for a postsynaptic leaky integrator, *Neuron*, **54**: 787-800, 2007.